



SPHERE THEOREM FOR MANIFOLDS WITH POSITIVE CURVATURE

Bazanfare Mahaman

► To cite this version:

Bazanfare Mahaman. SPHERE THEOREM FOR MANIFOLDS WITH POSITIVE CURVATURE. Glasgow Mathematical Journal, 2006, 48 (1), pp.37-40. hal-00002180v2

HAL Id: hal-00002180

<https://hal.science/hal-00002180v2>

Submitted on 15 Jul 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SPHERE THEOREM FOR MANIFOLDS WITH POSITIVE CURVATURE

BAZANFARÉ MAHAMAN

ABSTRACT. In this paper, we prove that, for any integer $n \geq 2$, there exists an $\epsilon_n \geq 0$ so that if M is an n -dimensional complete manifold with sectional curvature $K_M \geq 1$ and if M has conjugate radius bigger than $\frac{\pi}{2}$ and contains a geodesic loop of length $2(\pi - \epsilon_n)$, then M is diffeomorphic to the Euclidian unit sphere S^n .

1. INTRODUCTION

One of the fundamental problems in Riemannian geometry is to determine the relation between the topology and the geometry of a Riemannian manifold. In this way the Toponogov's theorem and the critical point theory play an important rule. Let M be a complete Riemannian manifold and fix a point p in M and define $d_p(x) = d(p, x)$. A point $q \neq p$ is called a critical point of d_p or simply of the point p if, for any nonzero vector $v \in T_q M$, there exists a minimal geodesic γ joining q to p such that the angle $(v, \gamma'(0)) \leq \frac{\pi}{2}$. Suppose M is an n -dimensional complete Riemannian manifold with sectional curvature $K_M \geq 1$. By Myers' theorem the diameter of M is bounded from above by π . In [Ch] Cheng showed that the maximal value π is attained if and only if M is isometric to the standard sphere. It was proved by Grove and Shiohama [GS] that if $K_M \geq 1$ and the diameter of M $diam(M) > \frac{\pi}{2}$ then M is homeomorphic to a sphere.

Hence the problem of removing homeomorphism to diffeomorphism or finding conditions to guarantee the diffeomorphism is of particular interest. In [Xi3] C. Xia showed that if $K_M \geq 1$ and the conjugate radius of M $\rho(M) > \pi/2$ and if M contains a geodesic loop of length 2π then M is isometric to $S^n(1)$.

1.1. Definition. *Let M be an n -dimensional Riemannian manifold and p be a point in M . Let $Conj(p)$ denote the set of first conjugate points to p on all geodesics issuing from p . The conjugate radius $\rho(p)$ of M*

at p is defined as

$$\rho(p) = d(p, \text{Conj}(p)) \quad \text{if } \text{Conj}(p) \neq \emptyset$$

and

$$\rho(p) = +\infty \quad \text{if } \text{Conj}(p) = \emptyset$$

Then the conjugate radius of M is:

$$\rho(M) = \inf_{x \in M} \rho(x).$$

Many interesting results have been proved by using the critical points theory and Toponogov's theorem [C], [GS], [Pe], [S], [Sh], [SS], [Xi1], [Xi2], [Xi3]. In otherewise J. Cheeger and T. Colding in [CC] have proven the following

Theorem A. *There exists a number, $\epsilon(n) > 0$, depending only on the integer n such that, for any two Riemannian manifolds Z_1, Z_2 , if $d_{GH}(Z_1, Z_2) < \epsilon(n)$, then Z_1 and Z_2 are diffeomorphic. where $d_{GH}(Z_1, Z_2)$ denote the Gromov-Hausdroff distance.*

The purpose of this paper is to prove the following :

1.2. Theorem. *For any $n \geq 2$ there exists a positif constant $\epsilon(n)$ depending only on n such that for any $\epsilon \leq \epsilon(n)$, if M is an n -dimensional complete connected Riemannian manifold with sectional curvature $K_M \geq 1$ and conjugate radius $\rho(M) > \frac{\pi}{2}$ and if M contains a geodesic loop of length $2(\pi - \epsilon)$ then M is diffeomorphic to an n -dimensional unit sphere $S^n(1)$.*

2. PROOF

Since $K_M \geq 1$, M is compact. Let $i(M)$ denote the injectivity radius of M .

By definition we have

$$i(M) = \inf_{x \in M} d(x, C(x)).$$

Since M is compact and the function $x \mapsto d(x, C(x))$ is continuous, there exists $p \in M$ such that $i(M) = d(p, C(p))$. Since $C(p)$ is compact there exists $q \in C(p)$ such that $i(M) = d(p, q) = d(p, C(p))$. Then

a) either there exists a minimal geodesic σ joining p to q such that q is a conjugate point of p or

b) there exists two minimal geodesics σ_1 and σ_2 joining p to q such that $\sigma'_1(l) = -\sigma'_2(l)$, $l = d(p, q)$. See [C].

2.1. Lemma. *Let M be an n -dimensional complete, connected Riemannian manifold with sectional curvature $K_M \geq 1$ and the conjugate radius $\rho(M) > \frac{\pi}{2}$, then $i(M) > \frac{\pi}{2}$.*

Proof of the lemma

If $a)$ holds, then $i(M) = d(p, q) > \pi/2$.

Suppose $b)$ holds. Since $q \in C(p)$, we have $p \in C(q)$ and consequently $d(p, q) = d(q, C(q))$. This implies that $\sigma'_1(0) = -\sigma'_2(0)$. Set $D(x) = \max_{y \in M} d(x, y)$. Then

$$D(x) \geq \max_{y \in C(x)} d(x, y) \geq \rho(M) > \pi/2.$$

Since M is compact, there exist a point $y \in M$ such that $D(x) = d(x, y) > \pi/2$ and y is the unique farthest point and the critical one for the distance $d(x, \cdot)$. Set $A(x) = y$; thus we define a continuous map $A : M \mapsto M$ (see [Xi3]). By the Berger-Klingenberg theorem, M is homeomorphic to the unit sphere $S^n(1)$ and since $A(x) \neq x$ for all $x \in M$ the Brouwer fixed point theorem sets that the degree of A is $(-1)^{n+1}$ and consequently A is surjective. Let $r \in M$ the point so that $p = A(r)$. Hence $d(p, r) > \pi/2$. If $r = q$ then $i(M) = d(p, q) > \pi/2$ otherwise there exists a minimal geodesic from q to r , σ_3 such that

$$\angle(\sigma'_3(0), -\sigma'_1(l)) \leq \pi/2 \quad \text{or} \quad \angle(\sigma'_3(0), -\sigma'_2(l)) \leq \pi/2.$$

Suppose $\angle(\sigma'_3(0), -\sigma'_1(l)) \leq \pi/2$. Applying the Toponogov's theorem [T] to the hinge (σ_1, σ_3) , we have:

$$\cos d(p, r) \geq \cos d(p, q) \cos d(q, r) + \sin d(p, q) \sin d(q, r) \cos \angle(\sigma'_3(0), -\sigma'_1(l))$$

$$(1) \quad \geq \cos d(p, q) \cos d(q, r).$$

Since r is far from p in the sense that $d(p, r) > \pi/2$ then r is near to q i.e $d(q, r) < \pi/2$ and from (1) we have

$$\cos d(p, q) < 0$$

and consequently

$$i(M) = d(p, q) > \pi/2$$

which proves the lemma.

2.2. Lemma. *Let M be a complete connected n -dimensional Riemannian manifold with sectional curvature $K_M \geq 1$ and conjugate radius $\rho(M) > \frac{\pi}{2}$. If M contains a geodesic loop of length at least $2(\pi - \epsilon)$ then $\text{diam}(M) \geq \pi - \tau(\epsilon)$ where $\tau(\epsilon) \mapsto 0$ when $\epsilon \mapsto 0$.*

Proof

Since $i(M) > \pi/2$ then there exist $\delta > 0$ such that $i(M) > \pi/2 + \delta$. Let γ be a loop with length $2\pi - 2\epsilon$. Let $x = \gamma(0) = \gamma(2\pi - 2\epsilon)$, $y = \gamma(\pi/2 + \delta)$, $m = \gamma(\pi - \epsilon)$ and $z = \gamma(\frac{3(\pi-\epsilon)}{2} - \delta)$

Let

$$\gamma_1 = \gamma/[0, \frac{\pi}{2} + \delta], \quad \gamma_2 = \gamma/[\frac{\pi}{2} + \delta, \pi - \epsilon] \quad \gamma_3 = \gamma/[\pi - \epsilon, \frac{3(\pi-\epsilon)}{2} - \delta]$$

and $\gamma_4 = \gamma/[\frac{3(\pi-\epsilon)}{2} - \delta, 2\pi - 2\epsilon]$.

Then the geodesics γ_i are minimal. Let σ be a minimal geodesic joining m and x .

We claim that $L(\sigma) \geq \pi - \tau(\epsilon)$.

Set $\alpha = \angle(\sigma'(0), -\gamma'(\pi - \epsilon))$ and $\beta = \angle(\sigma'(0), \gamma'(\pi - \epsilon))$. Applying the Toponogov's theorem to the triangles $(\gamma_1, \gamma_2, \sigma)$ and $(\gamma_3, \gamma_4, \sigma)$ respectively, one can take two triangles $(\bar{\gamma}_1, \bar{\gamma}_2, \bar{\sigma})$ and $(\bar{\gamma}_3, \bar{\gamma}_4, \bar{\sigma})$ on the unit sphere $S^2(1)$ with vertices $\bar{x}, \bar{y}, \bar{m}$ and $\bar{x}, \bar{z}, \bar{m}$ respectively satisfying:

$$L(\bar{\gamma}_i) = L(\gamma_i), \quad i = 1, 2, 3, 4; L(\bar{\sigma}) = L(\sigma);$$

hence $\bar{\alpha} \leq \alpha, \bar{\beta} \leq \beta$ where $\bar{\alpha}$ and $\bar{\beta}$ are the angles at \bar{m} of the triangles $(\bar{\gamma}_1, \bar{\gamma}_2, \bar{\sigma})$ and $(\bar{\gamma}_3, \bar{\gamma}_4, \bar{\sigma})$ respectively. We have: $\alpha \leq \pi/2$ or $\beta \leq \pi/2$. Suppose, without lost the generality, that $\alpha \leq \pi/2$. Let \bar{x}' be the antipodal point of \bar{x} on the sphere $S^2(1)$ and $\bar{\sigma}_1$ the minimal geodesic from \bar{m} to \bar{x}' . If $\bar{\alpha}'$ and $\bar{\beta}'$ are the angles at \bar{m} , of triangles $(\bar{y}, \bar{m}, \bar{x}')$ and $(\bar{z}, \bar{m}, \bar{x}')$ respectively then we have:

$$\begin{aligned} d((\bar{y}, \bar{x}')) &= \pi - d(\bar{y}, \bar{x}) \\ (2) \quad &= \pi - d(x, y) = \frac{\pi}{2} - \delta \end{aligned}$$

Hence, using the trigonometric law on the triangle $(\bar{y}, \bar{m}, \bar{x}')$ we have:

$$\begin{aligned} \sin d(\bar{m}, \bar{y}) \sin d(\bar{m}, \bar{x}') \cos \bar{\alpha}' &= \cos d(\bar{y}, \bar{x}') - \cos d(\bar{m}, \bar{y}) \cdot \cos d(\bar{m}, \bar{x}') \\ &= \cos\left(\frac{\pi}{2} - \delta\right) - \cos\left(\frac{\pi}{2} - \delta - \epsilon\right) \cdot \cos d(\bar{m}, \bar{x}') \\ &= \sin(\delta) - \sin(\delta + \epsilon) \cos d(\bar{m}, \bar{x}') \leq 0 \end{aligned}$$

which means that

$$\cos d(\bar{m}, \bar{x}') \geq \frac{\sin(\delta)}{\sin(\delta + \epsilon)}.$$

It follows that

$$L(\sigma) = d(\bar{m}, \bar{x}') \leq \cos^{-1} \left(\frac{\sin(\delta)}{\sin(\delta + \epsilon)} \right) = \tau(\epsilon)$$

with $\tau(\epsilon) \mapsto 0$ when $\epsilon \mapsto 0$.

Hence

$$d(m, x) = d(\bar{m}, \bar{x}) \geq \pi - \tau(\epsilon).$$

2.3. Lemma. *Let M be a complete connected n -dimensional Riemannian manifold with sectional curvature $K_M \geq 1$ and $\text{diam}(M) \geq \pi - \epsilon$, then for all $x \in M$ there exists a point x' such that $d(x, x') \geq \pi - \Gamma(\epsilon)$ hence $\text{Rad}(M) \geq \pi - \Gamma(\epsilon)$ with $\Gamma(\epsilon) \mapsto 0$ when $\epsilon \mapsto 0$ where*

$$\text{Rad}(M) = \min_{x \in M} \max_{y \in M} d(x, y)$$

Proof

Since M is compact, its injectivity radius $i(M)$ is positive and set r_0 a positive number not larger than $i(M)$. Let p, q be two points in M such that $d(p, q) = \text{diam}(M) \geq \pi - \epsilon$. Let $x \in M$ and suppose $x \neq p$ and $x \neq q$. Consider the triangle (p, q, x) and let y be a point of the segment $[q, x]$ such that $d(q, y) = \frac{r}{2}$ with $r < r_0$ and $d(p, x) > 2r$ $d(q, x) > 2r$. For any point $s \in B(y, r)$ the function

$$z \mapsto e_{yz}(s) = d(y, s) + d(z, s) - d(y, z)$$

is continuous and if z is on the prolongation of the geodesic joining y to s we have: $e_{yz}(s) = 0$ (this is possible since $r < r_0$).

Hence the function $z \mapsto e_{yz}(q)$ is continuous on the sphere $S(y, r)$, and consequently there exists $z \in S(y, r)$ such that

$$(3) \quad e_{yz}(q) = d(y, q) + d(z, q) - d(y, z) < \epsilon$$

For any point $v \in M$ we have:

$$d(p, v) + d(q, v) + d(p, q) \leq 2\pi$$

hence

$$(4) \quad |d(p, v) + d(q, v) - \pi| \leq \epsilon.$$

Let \bar{p}' be the antipodal point of \bar{p} on the sphere S^2 .

$$d(\bar{p}, \bar{q}) = d(p, q);$$

$$d(\bar{z}, \bar{p}') = \pi - d(\bar{z}, \bar{p}) = \pi - d(p, z).$$

We have

$$|d(\bar{q}, \bar{y}) - d(\bar{y}, \bar{p}')| = |d(\bar{q}, \bar{y}) - \pi + d(\bar{p}, \bar{y})| < \epsilon.$$

In the same way, we have:

$$|d(\bar{q}, \bar{z}) - \pi + d(\bar{p}, \bar{z})| = |d(q, z) - \pi + d(p, z)| < \epsilon$$

Hence

$$(5) \quad \begin{aligned} d(\bar{y}, \bar{z}) &= d(y, z) + d(q, y) + d(q, z) - \epsilon \geq \pi - \epsilon - d(p, y) + d(q, z) - \epsilon \\ &\geq \pi - d(\bar{p}, \bar{y}) + d(\bar{q}, \bar{z}) - 2\epsilon \geq d(\bar{p}', \bar{y}) + d(\bar{q}, \bar{z}) - 2\epsilon \end{aligned}$$

$$(6) \quad \geq d(\bar{p}', \bar{y}) + d(\bar{p}', \bar{z}) - 3\epsilon$$

Thus $d(\bar{p}', \bar{y}) > \frac{r}{2} - \frac{3}{2}\epsilon$ and $d(\bar{p}', \bar{z}) > \frac{r}{2} - \frac{3}{2}\epsilon$.

Suppose $d(\bar{p}', \bar{y}) \leq \frac{r}{2} - \frac{3}{2}\epsilon$ then $d(\bar{p}, \bar{y}) \geq \pi - \frac{r}{2} + \frac{3}{2}\epsilon$ which contradicts (4).

Let

$$\bar{l}'_1 = d(\bar{p}', \bar{y}); \bar{l}'_2 = d(\bar{p}', \bar{z}) \text{ and } \bar{l}'_0 = d(\bar{z}, \bar{y}).$$

Applying the Topogonov's theorem to the triangle $(\bar{y}, \bar{p}', \bar{z})$, we have:

$$\sin \bar{l}'_1 \sin \bar{l}'_2 \cos \angle \bar{p}' = \cos \bar{l}'_0 - \cos \bar{l}'_1 \cos \bar{l}'_2$$

From inequality (6) we get:

$$\begin{aligned} \sin \bar{l}'_1 \sin \bar{l}'_2 \cos \angle \bar{p}' &< \cos(\bar{l}'_1 + \bar{l}'_2 - 3\epsilon) - \cos \bar{l}'_1 \cos \bar{l}'_2 \\ &< -\cos \bar{l}'_1 \cos \bar{l}'_2 (1 - \cos 3\epsilon) - \sin \bar{l}'_1 \sin \bar{l}'_2 \cos 3\epsilon - \sin(\bar{l}'_1 + \bar{l}'_2) \sin 3\epsilon \end{aligned}$$

Hence

$$\cos \angle \bar{p}' < -\cos 3\epsilon + (\cot g \bar{l}'_2 + \cot g \bar{l}'_1) \sin 3\epsilon - \cot g \bar{l}'_1 \cdot \cot g \bar{l}'_2 (1 - \cos 3\epsilon).$$

Thus $\angle \bar{p}' > \pi - \tau_1(\epsilon)$ with

$$\tau_1(\epsilon) = \cos^{-1} \left(\cos 3\epsilon - (\cot g \bar{l}'_2 + \cot g \bar{l}'_1) \sin 3\epsilon + \cot g \bar{l}'_1 \cdot \cot g \bar{l}'_2 (1 - \cos 3\epsilon) \right)$$

and $\tau_1(\epsilon) \mapsto 0$ as $\epsilon \mapsto 0$.

The trigonometric law on the sphere shows that the angle at \bar{p}' of the triangle $(\bar{y}, \bar{p}', \bar{z})$ is equal to the angle at \bar{p} of the triangle $(\bar{y}, \bar{p}, \bar{z})$ which is not bigger than the angle at p of the triangle (y, p, z) in M .

Hence $\angle(y, p, z) > \pi - \tau_1(\epsilon)$.

By applying the relation (4) to x and y we have:

$$(7) \quad -\epsilon \leq d(p, x) + d(q, x) - \pi \leq \epsilon,$$

$$(8) \quad -\epsilon \leq d(p, y) + d(q, y) - \pi \leq \epsilon.$$

Since $y \in [q, r]$, we conclude from (7) and (8) that $e_{py}(x) < 2\epsilon$, which shows that the angle at x of the triangle (p, x, y) is close to π and consequently its angle at p is small. there exists

$\tau_2(\epsilon)$ such that $\angle p \leq \tau_2(\epsilon)$ where $\tau_2(\epsilon) \mapsto 0$ as $\epsilon \mapsto 0$.

Take r small enough; then $d(p, x) < d(p, y)$; otherwise $d(p, x)$ is close to π and we conclude by taking $x' = p$.

Let $\tilde{x} \in [p, y]$ such that $d(p, \tilde{x}) = d(p, x)$; then

$$(9) \quad d(x, \tilde{x}) \leq \pi \angle(y, p, x) = \tau_3(\epsilon).$$

Since

$$d(p, z) + d(z, q) \geq d(p, q) \geq \pi - \epsilon$$

and

$$d(y, z) = r > d(q, y) + d(q, z) - \epsilon$$

we have:

if $d(p, x) \leq d(q, z) + \epsilon$ then $r < \frac{2}{3}\epsilon$ and it suffices to take $x' = q$ and $\Gamma(\epsilon) \leq \frac{7}{3}\epsilon$; if $d(p, x) > d(q, z) + \epsilon$ then

$$d(p, z) \geq \pi - d(q, z) - \epsilon > \pi - d(p, x)$$

hence there exists a point $x' \in [p, z]$ such that

$$d(p, x') = \pi - d(p, x).$$

It suffices to show that $d(x', \tilde{x}) \geq \pi - \tau_4(\epsilon)$.

Applying the Toponogov theorem to the triangle (\tilde{x}, p, x') , we get

$$\begin{aligned} \cos d(x', \tilde{x}) &= \cos d(p, x') \cos d(p, \tilde{x}) + \sin d(p, x') \sin d(p, \tilde{x}) \cos \angle p \\ &= -\cos^2 d(p, \tilde{x}) + \sin^2 d(p, \tilde{x}) \cos \angle p \end{aligned}$$

Since $x' \in [p, z]$ and $\tilde{x} \in [p, y]$ the angle at \bar{p} of the triangle $(\bar{y}, \bar{p}, \bar{z})$ is less or equal to the angle at \bar{p} of the triangle $(\tilde{x}, \bar{p}, \bar{x}')$ which is not bigger than the angle at p of triangle (\tilde{x}, p, x') in M .

$$\angle(\tilde{x}, p, x') \geq \angle(\tilde{x}, \bar{p}, \bar{x}') \geq \angle(\bar{y}, \bar{p}, \bar{z}) \geq \pi - \tau_4(\epsilon).$$

Hence

$$\begin{aligned} \cos d(x', \tilde{x}) &\leq -\cos^2 d(p, \tilde{x}) + \sin^2 d(p, \tilde{x}) \cos(\pi - \tau_4(\epsilon)) \\ &= -\cos(\tau_4(\epsilon)) - \cos^2 d(p, x) (1 - \cos(\tau_4(\epsilon))) \leq -\cos(\tau_4(\epsilon)) \\ &\Rightarrow d(\tilde{x}, x') > \pi - \tau_4(\epsilon). \end{aligned}$$

From the triangle inequality and the inequality (9) we have

$$d(x, x') \geq d(\tilde{x}, x') - d(\tilde{x}, x) \geq \pi - \tau_5$$

Thus, lemma 2.3 follows.

In [Gr] M. Gromov generalized the classic notion of Hausdorff distance between two compact subsets of the same metric space. He considered the set of compact Riemannian manifolds as a subset of the set of all compact metric spaces.

2.4. Definitions. 1) Let X, Y be two metric spaces; a map $f : X \rightarrow Y$ is said to be an ϵ -approximation if the image set $f(X)$ is ϵ -dense in Y and, for any $x, y \in X$, $|d(f(x), f(y)) - d(x, y)| < \epsilon$.

2) The Gromov-Hausdorff distance $d_{GH}(X, Y)$ between X and Y is the infimum of values of $\epsilon > 0$ such that there exist ϵ -approximations $f : X \rightarrow Y$ and $g : Y \rightarrow X$. In [Co1] and [Co2] Colding showed the two equivalent conditions:

- 1) $Rad(M) \geq \pi - \epsilon$
- 2) $d_{GH}(M, S^n(1)) \leq \tau_5(\epsilon)$ with $\tau_5(\epsilon) \mapsto 0$ as $\epsilon \mapsto 0$.

Then theorem 1.2 follows from these conditions and the theorem A.

3. REFERENCES

- [C] J.Cheeger, Critical points of distance functions and applications to geometry, Lectures notes 1504 (1991) 1-38.
- [CC] J.Cheeger and T. Colding, On the structure of spaces with Ricci curvature bounded from below, J. Diff. Geom. 46 (1997) 406-480.
- [Ch] S.Y.Cheng, Eigenvalue Comparison Theorem and Geometric Applications Math.Z.143(1975) p. 289-297.
- [Co1] T.H. Colding, Shape of manifolds with positive Ricci curvature, Invent. Math. 124 (1996), 175-191
- [Co2] T.H. Colding, Large manifolds with positive Ricci curvature, Invent. Math. 124 (1996), 193-214
- [Gr] M.Gromov, Structures métriques pour les variétés Riemanniennes, Rédigé par J.Lafontaine et P.Pansu, cedic 1981.
- [GS] K. Grove and K. Shiohama, A generalized sphere theorem, Ann. Math. 106 (1977), 201-211.
- [Pe] P. Petersen, Comparison geometry problem list, Riemannian geometry (Waterloo, ON, (1993) 87-115, fields Inst. Monogr., 4, Amer. Math. Soc., Providence, RI 1996.
- [S] K. Shiohama, A sphere theorem for manifolds of positive Ricci curvature, Trans. Amer. Math. Soc. 275 n.2 (1983) 811-819.
- [Sh] Z. Shen, Complete manifolds with nonnegative Ricci curvature and large volume growth, Invent. Math. 125 (1996) 393-404.
- [SS] J. Sha and Z. Shen, Complete manifolds with nonnegative curvature and quadratically nonnegative curved infinity, Amer. J. Math. 119 (1997) 1399-1404.
- [T] V.A.Toponogov, Computation of length of a closed geodesic on convex surface, Dokl. akad. Nank SSSR 124 (1959) 282-284.
- [Xi1] C.Y. Xia, Large volume growth and topology of open manifolds, Math. Z. 239 (2002) 515-526.
- [Xi2] " " , Complete manifolds with sectional curvature bounded below

and large volume growth, Bull. London Math. Soc. 34 (2002) 229-235.
 [Xi3] ” ” , Some applications of critical point theory of distance functions on Riemannian manifolds, Compos.math.132, n.1(2002)pp 49-55.

Bazanfaré Mahaman
 Université Abdou Moumouni
 Département de Mathématiques et Informatique Niamey- Niger
 E-mail: bmahaman @yahoo.fr